UNIVERSITY OF AGRICULTURAL SCIENCE, BENGALURU
GRAMIN KRISHI MAUSAM SEWA(GKMS)
AMFU OF IMD, BENGALURU

AGROMET-ADVISORY BULLETIN Date: **28.05.2019**

Issued jointly by, UAS, Bengaluru & Indian Meteorological Department

Pest forecast (Valid from 29th May 2019 to 02nd June 2019)

Pest Advisory:

Crop	Stage/ Condition	Pest and Disease	Agro advisories
Agriculture crop			
Kharif crops	➢ Summer ploughing and FYM application in field to manage soil borne disease ➢ In mango, due to heavy downpour of rain in thunderstorm fruit drop have been observed in mango consequently fruit fly damage is also noticed. suitable control measure by spraying Carbaryl 4 g/litre of water		

The preventive control practices for paddy pests and disease problems

1. **Practice good cleaning of equipment and field between seasons**
2. **Use clean seeds and resistant varieties**
3. **Plant at the same time as your neighbors**
   - Planting at the same time (or within a 2-week window) as the neighboring fields can help to minimize insect, disease, bird, and rat pressure on individual fields
4. **Do not over apply fertilizer**
   - Application of recommended dose of fertilizers
5. **Encourage natural pest enemies**
6. **Do not apply pesticide within 40 days of planting**
7. **Properly store grain**
   - Store grain at moisture content below 13-14%, preferably in an airtight container
   - Do not store grain for more than 6 months